

# Adaptation of Internet Addiction Scale in Azerbaijani Language: A Validity-reliability and Prevalence Study

Melek Kerimova<sup>1</sup>, Selim Gunuc<sup>2</sup>

<sup>1</sup>School of Humanities and Social Sciences, Khazar University, Baku, Azerbaijan

<sup>2</sup>Department of Computer Education and Instructional Technologies, Yuzuncu Yil University, Turkey

Correspondence: Selim Gunuc, Department of Computer Education and Instructional Technologies, Yuzuncu Yil University, Turkey

Received: July 30, 2015 Accepted: September 6, 2015 Online Published: September 25, 2015

doi:10.11114/jets.v4i1.1105

URL: <http://dx.doi.org/10.11114/jets.v4i1.1105>

## Abstract

The purpose of the present paper was to adapt Gunuc and Kayri's (2010) *Internet Addiction Scale*, with show validity and reliability for many various sampling groups, into the Azerbaijani language. Another objective of the study is to determine the prevalence of Internet addiction among Azerbaijani adolescents and youth, which preemptively requires adaptation of the Internet Addiction Scale into the Azerbaijani language. Samples in the study have been selected separately for both purposes (validity and prevalence). Data were collected from students attending Khazar University in Azerbaijan in 2014-2015, first, in order to conduct Confirmatory Factor Analysis of the scale and second, to determine the prevalence of Internet addiction among youth. Data was obtained from 309 students for the first purpose and from 836 students for the second one. Students in various majors within the research sample were stratified through stratified sampling and students in these majors were randomly selected. The four-factor structure of the scale was tested through CFA. The Cronbach's Alpha ( $\alpha$ ) internal consistency coefficient of the scale was calculated as  $\alpha=.926$ . Reliability coefficient of 4 sub-factors, regarding the scale, was found as follows:  $\alpha=.844$  for *Withdrawal*,  $\alpha=.821$  for *Controlling difficulty*,  $\alpha=.829$  for *Disorder in Functionality* and  $\alpha=.792$  for *Social Isolation*. The prevalence of Internet addiction among youth in Azerbaijan was determined. The study found that 51.9% were not Internet addicts, 40.6% were moderate Internet addicts, and 7.5% were Internet addicts. These findings show that the rate of Internet addicts, particularly in the categories of moderate Internet addicts and Internet addicts (48.1%), is not small enough to ignore.

**Keywords:** Internet addiction, scale, Azerbaijan, adolescent, prevalence, validity

## 1. Introduction

Based on the data presented by Miniwatts Marketing Group (2014), it is stated that the global population is 7.264.623.793 and globally 3.079.339.857 people have access to the Internet. This number represents the 42.4% of the world population. An increase of 676.3% is mentioned around the world between 2000 and 2013. Regarding the number of Internet users, China and the USA take the first places whereas Turkey ranks 18<sup>th</sup>. In 2000, Turkey had 2.000.000 Internet users but in 2013 this number went up to 37.748.969 (<http://www.internetworldstats.com/stats.htm>).

The Internet in Azerbaijan was first used at the Institute of Information Technologies in National Sciences Academy in 1991. First email service was offered within this institute. The first Internet connection in Azerbaijan in 1994 was provided at Management Branch of Azerbaijan National Sciences Academy, with support from British Petroleum and Turkish Government and Middle East Technical University participation, through Turksat satellite. In 1994, "www.ab.az" website for the first time was created and in 1997, the first government website "www.prezident.az" was built. According to the data presented by Azerbaijan State Statistical Committee (2013), the nationwide rate of households with a computer is 53.2% and Internet access is 71.6%. Based on the data by "APA-ECONOMICS" (2010), the number of Internet users in Azerbaijan accounts for 44.4% of the population. This approximately adds up to 3.7 million users. For Azerbaijan, a country with a population of 9.5 million, this number is very crucial (<http://www.stat.gov.az/-12.05.2014>). Therefore, Internet addiction, as the unwanted face of Internet use, is a concern.

Although the increase in the number of Internet users gives a positive impression based on considerations about possibilities and benefits of Internet use, the status of Internet addiction invokes anxiety and concern. Internet addiction is considered within behavioral addictions such as TV addiction, eating addiction, and shopping addiction. Although, in

types of substance addiction, concrete symptoms and damages regarding the addicted person's health can be mentioned, it is quite hard to talk about overall damages and, therefore, to define the line between the Internet use and addiction.

It is observed that the concept has been used in various forms within the relevant academic literature. Dr. Ivan Goldberg (1996) was the first to use it in this sense as *Internet addiction* on international level and it is seen that it was used as *Internet dependency*. Young defined the concept of Internet addiction, based on *pathologic gambling criteria* of DSM IV. Internet addiction is considered a new psychiatric disorder within DSM-5 (2013). Researchers following Young and Goldberg who suggest using the concept as *pathological Internet use* attract attention. Actually, it may be stated that all concepts used share a common ground. It can be said that the most highlighted notion among all of the concepts about extreme and problematic Internet use is related to wasting most of an individual's time on the Internet. For Internet addiction, although time is a criterion, the purpose of using time is important. Individuals who are not Internet-addicted use it for informational, search, and other purposes. However, Internet-addicted individuals are online for long hours and cannot be away from the Internet.

According to international literature, the spread of Internet addiction among adolescents in several countries is as follows: 10.1% in Turkey (Gunuc & Kayri, 2010), 5.8% in Poland (Zboralski et al., 2009), 5.8% in Italy (Poli & Agrimi, 2012), 13.5% in China (Wu et al., 2013), 4.6% in Romania (Durkee et al., 2012), 17.7% in Romania (Tsitsika et al., 2014), 5.1% in Germany (Wölfling & Muller, 2010), 10.6% in Germany (Tsitsika et al., 2014), 4.4% in Europe (Durkee et al., 2012) and 13.9% in Europe (Tsitsika et al., 2014). It could be stated that these differences from one society to another result from a number of factors such as culture, access to technology, age, parental education, level of income, different measurement tools and different measurement techniques, time of research conducted and selection of research samples (Durkee et al., 2012; Tsitsika et al., 2014).

When distinguishing between an Internet-addicted person and others, many tests, scales, questionnaires, and surveys are used around the world. Today, it is important to accurately define the Internet addiction. In this sense, the need for the scales is on the increase and it can presently be said that the scales are among important economical, practical, and reliable methods of measurement. Although no scales related to Internet addiction is available in Azerbaijan, when relevant international studies are reviewed, it is observed that various measurement tools are used (Young, 1998; Caplan, 2002; Thatcher & Goolam, 2005; Caplan, 2010; Clark & Frith, 2005). Some scales, tests, and inventories of Internet addiction are primarily observed when literature about relevant measurement tools is reviewed.

An 8-question "Diagnostic Criteria for Internet Addiction" scale developed by Young in 1996 based on DSM-IV "Substance Addiction" criteria is considered the first among scales internationally utilized in relation to Internet addiction. Later, Young (1998) improved this scale (IAT), adding up to 20 items. This scale has been used with adaptations in countries such as China, Korea, Germany, and Turkey. "The Pathological Internet Use" scale developed by Morahan-Martin and Schumacher (2000) is a Likert-type scale with 13 items.

In addition, various measurement tools such as Caplan's (2002) 5-point Likert-type "Generalized Problematic Internet Use Scale" with 29 items, "Chinese Internet Addiction Inventory" with 31 items developed by Chen and colleagues (2003), "Internet Addiction Scale" with 31 items developed by Nichols and Nicki (2004), "Problematic Internet Use Questionnaire" with 20 items developed by Thatcher and Goolam (2005), "Internet Addiction Scale" with 20 items developed by Kim and colleagues (2006), and "Problematic Internet Use Questionnaire" with 18 items developed by Demetrovics and colleagues (2008) are available. Lastly, "Internet Addiction Scale" with 35 items, developed by Gunuc and Kayri (2010), used in many studies in Turkey, a country culturally close to Azerbaijan, ranks among salient measurement instruments.

As can be observed, prevalence of Internet addiction varies in different countries and the measures used vary as well. In this sense, measures with proven validity and reliability for various samples become more important. Yet, although many studies have been conducted in different countries, it is observed that Azerbaijan lacks empirical research in this field. In this sense, it attracts attention that both no measurement instrument in regards to Internet addiction and no empirical studies on the issue are available in Azerbaijan. Due to the cultural and linguistic similarities between Azerbaijan and Turkish societies, the purpose of this study is to adapt Gunuc and Kayri's (2010) *Internet Addiction Scale*, with show validity and reliability for many various sampling groups and ages (such as Gunuc, 2015; Gunuc, 2013; Gunuc, 2011; Gunuc & Dogan, 2013), into the Azerbaijani language. The study also aims to ascertain the prevalence of Internet addiction among Azerbaijani adolescents and youth, which preemptively requires adaptation of the *Internet Addiction Scale* into the Azerbaijani language.

The present research endeavor, as the first empirical study, qualifies as a pilot study in Azerbaijan to help accurately determine Internet addicts. In other words, the adapted scale is intended as a tool to help define the Internet addicts in Azerbaijan. In addition to clinical observations, the scale is expected to contribute toward defining the prevalence of Internet addicts in larger and different sample groups.

## 2. Method

### 2.1 Sample

The population for this research project consists of students attending Khazar University. The ages of the participants varied between 18 and 22. Data were collected in classrooms on pencil-and-paper basis in 2014-2015 academic years. Samples in the study have been selected separately for both purposes (validity and prevalence). Data were collected from students attending Azerbaijan Khazar University in 2014-2015, first, for the purpose of *Confirmatory Factor Analysis* of the scale (validity) and second, in order to determine the prevalence of Internet addiction. Students in various majors within the research sample were stratified through stratified sampling and students in these majors were randomly selected. Data were obtained from 309 students in the first purpose category and from 836 students in the second purpose category.

### 2.2 Data Collection Tools

“Personal Information Form” prepared by researchers and “Internet Addiction Scale” developed by Gunuc and Kayri (2010) as data collection tools were used in the present study. Permission to administer the data collection instruments among students was obtained from Khazar University president’s office and relevant explanation was provided to the students about how and where the research results would be used in order for them to provide accurate response.

#### 2.2.1 Internet Addiction Scale (IAS)

Because IAS was adapted into the Azerbaijani language, it was tested through its original factor structure CFA, without conducting exploratory factor analysis. IAS was developed in Turkish by Gunuc and Kayri (2010) via a study carried out with 754 adolescents. The average of participants was 15.8. The scale was made up of 35 items. In their study, the Cronbach’s Alpha internal consistency coefficient of the five-point Likert-type scale, rated as (1) *strongly disagree* and (5) *strongly agree*, was calculated by the researchers as .94. The Cronbach’s Alpha (a) reliability coefficients, regarding the four sub-factors of IAS, were calculated as  $a=.877$  for the sub-factor of Withdrawal, as  $a=.855$  for the sub-factor of Controlling Difficulty, as  $a=.827$  for the sub-factor of Disorder in Functionality and as  $a=.791$  for the sub-factor of Social Isolation. Higher scores received on the scale indicate Internet addiction.

According to Hambleton and Patsula (1999), just any two bilinguals will not suffice in adapting a scale; they must also be equipped with relevant field expertise. Based on the aforesaid, the original scale was translated into the Azerbaijani language by two bilinguals, of both Azerbaijani and Turkish languages, as experts of Psychological Counselling and Guidance and Cyber-Psychology. No significant differences were noticed between the expert translations. Later, as Hambleton and Patsula (1999) stated, the scale was translated into Turkish by the two experts and their consistency with original item structures was examined. Items on the original scale and the translated items in Turkish were compared on semantic, conceptual, idiomatic, and experiential criteria. Then, the translated scale was administered on 25 students attending various departments and revised based on feedback from the individuals, to complete the process.

### 2.3 Data Analysis for CFA

The goodness-of-fit indices for the model tested through CFA was determined with the help of  $\chi^2$  (Chi-Square Goodness-of-Fit), GFI (Goodness-of-Fit Index), AGFI (Adjusted Goodness-of-Fit Index), CFI (Comparative Fit Index), NFI (Normed Fit Index), NNFI (Not-Normed Fit Index), RMR (Root Mean Square Residuals), SRMR (Standardized Root Mean Square Residuals) and RMSEA (Root Mean Square Error of Approximation) indices. For the analysis of the data, the package programs of SPSS 18.0 and Lirsel 8.5 (Jöreskog & Sörbom, 2001) were used for descriptive statistics and confirmatory factor analysis, respectively.

### 2.4 Data Analysis for Prevalance

For prevalence, Two-Step Cluster Analysis was conducted to examine whether the participants were Internet addicts. Cluster analysis is a multivariate statistical technique to categorize individuals or objects in sub-classes or clusters depending on their similarities. The purpose is to gather individuals with similar characteristics in the same group considering a certain characteristic. For this purpose, the similarities or distances between units are used as a criterion (Everitt, 1980; Kaufman & Rousseeuw, 1990). In this study, the total scores obtained via IAS were used as the criterion for grouping. In this respect, Two-Step Cluster Analysis was conducted; the addicted and non-addicted adolescents were divided in groups; and the profile of the research sample was defined. For the analyses, the significance level was taken as .05.

## 3. Findings

### 3.1 Findings Regarding the Adaptation of Internet Addiction Scale

#### 3.1.1 Examining the Assumptions

For CFA, regarding the data collected from 309 participants found in the sample, z-scores for the univariate outliers and

Mahalanobis distances for the multivariate outliers were calculated (Huck, 2012; Kline, 2011). 6 students with outliers were excluded from the data set. As the values of both z-scores and Mahalanobis output for are all assumptions of multivariate analyses, were examined (Hutcheson & Sofroniou, 1999; Kline, 2011; Tabachnick & Fidell, 2007). The values of skewness (.252;  $\pm 1$ ) and kurtosis (-.178;  $\pm 1$ ) were found to be in acceptable range. In addition, when the histogram, Kolmogorov-Smirnov Normality Test (.200;  $p > .05$ ), P-P and Q-Q graphics for CFA were examined (Huck, 2012; Pallant, 2007; Kline, 2009), it was seen that the distributions were normal.

### 3.1.2 Confirmatory Factor Analysis (CFA)

The IAS structure was made up of 35 four-factor items. As the data set demonstrated a normal distribution, *Maximum Likelihood Method* as the parameter estimation method and *Covariance Matrix* as the data matrix were used in CFA. As can be seen in Table 1, as a result of CFA conducted, the outputs were examined, and primarily the *t*-values, factor loadings and error variances were evaluated.

Table 1. Item Statistics Regarding the CFA Findings

Item	<i>t</i>	Factor loading	Error variance	R <sup>2</sup>	Item-Total scale cor. (r)	<i>X</i>	Sd
<b>Withdrawal</b>		<b><math>\alpha = .844</math></b>					
M1	7.91*	.46	.79	.22	.448**	2.50	1.280
M2	9.03*	.51	.74	.30	.484**	2.95	1.342
M3	7.55*	.44	.81	.23	.413**	3.57	1.220
M4	9.52*	.54	.71	.28	.442**	2.31	1.144
M5	11.20*	.61	.62	.37	.525**	2.82	1.302
M6	13.98*	.73	.47	.53	.608**	2.57	1.234
M7	12.87*	.68	.53	.46	.603**	2.34	1.276
M8	9.51*	.54	.71	.29	.444**	2.28	1.197
M9	12.07*	.65	.58	.42	.618**	2.25	1.236
M10	13.67*	.72	.49	.51	.629**	2.58	1.351
M11	10.20*	.57	.68	.32	.508**	2.82	1.357
<b>Controlling Difficulty</b>		<b><math>\alpha = .821</math></b>					
M12	10.24*	.57	.68	.33	.567**	2.52	1.260
M13	9.36*	.53	.72	.28	.538**	2.62	1.402
M14	10.52*	.58	.66	.34	.533**	2.09	1.035
M15	9.92*	.55	.69	.31	.555**	2.85	1.330
M16	9.91*	.55	.69	.30	.521**	1.98	1.079
M17	10.98*	.60	.64	.36	.517**	2.10	1.220
M18	11.12*	.61	.63	.37	.536**	1.79	1.056
M19	11.20*	.61	.63	.38	.534**	2.09	1.145
M20	12.40*	.66	.56	.44	.584**	1.91	1.079
M21	11.88*	.64	.59	.42	.590**	2.39	1.354
<b>Disorder in Functionality</b>		<b><math>\alpha = .829</math></b>					
M22	12.13*	.65	.57	.41	.530**	1.95	1.168
M23	12.18*	.66	.57	.42	.546**	1.62	.871
M24	13.25*	.70	.51	.55	.632**	2.08	1.181
M25	12.80*	.69	.53	.53	.637**	2.34	1.315
M26	10.89*	.60	.64	.35	.567**	2.31	1.235
M27	12.89*	.69	.53	.48	.589**	2.13	1.195
M28	12.18*	.66	.57	.41	.531**	1.70	.941
<b>Social Isolation</b>		<b><math>\alpha = .792</math></b>					
M29	12.40*	.67	.55	.43	.454**	1.44	.673
M30	12.98*	.70	.51	.46	.443**	1.46	.824
M31	10.58*	.59	.65	.34	.320**	1.52	.881
M32	11.27*	.63	.61	.38	.461**	1.68	.931
M33	12.86*	.69	.52	.54	.603**	1.70	.955
M34	9.49*	.55	.70	.37	.590**	2.05	1.212
M35	12.12*	.66	.56	.42	.495**	1.86	1.147
Total-Scale Reliability		$\alpha = .926$					

\* $p < .01$ ; \*\* $p < .001$

As can be seen in Table 1, the *t*-value for each item was higher than +2.58, and the error variance was lower than .90. The *t*-value for each indicator in the scale is suggested to be out of the range of +2.58 ( $p < .01$ ) (Kline, 2011; Raykov & Marcoulides, 2006; Tabachnick & Fidell, 2007). Accordingly, it was seen that the items had a high level of *t*-value and that the error variance was not much high. The path diagram regarding the model can be seen in Figure 1.




Figure 1. Standardized Path Diagram

When the fit indices of the model were taken into consideration, the  $p$  level for the  $\chi^2$  value was examined. If this value is  $p > .05$ , then it shows good fit. However, as this value is likely to be significant ( $p < .05$ ) for large sizes of samples, it is suggested that the ratio of  $\chi^2/df$  and other fit indices should be evaluated (Tabachnick & Fidell, 2007). The Chi-Square value was calculated as  $\chi^2 = 1115.99$  and degree of freedom as  $df = 551$ . If this value is lower than 3, then, it shows good fit (Kline, 2011; Tabachnick & Fidell, 2007). In this respect, the ratio of  $\chi^2/df$  ( $1115.99/551$ ) was calculated as 2.03. In addition, the other fit indices are presented in Table 2 and evaluated in line with the related literature.

Table 2. Evaluation of Fit Indices Regarding CFA

Index	Sample statistic	Perfect fit	Good fit	Decision	Rationale
$\chi^2/df$	2.03	$\chi^2/df \leq 2$	$\chi^2/df \leq 3$	Perfect fit	(Kline, 2011)
RMSEA	.058	RMSEA $\leq$ .05	RMSEA $\leq$ .08	Good fit	Hooper, Coughlan & Mullen (2008)
RMR	.078	RMR $\leq$ .05	RMR $\leq$ .08	Good fit	Brown (2006) Hu & Bentler (1999)
SRMR	.064	SRMR $\leq$ .05	SRMR $\leq$ .08	Good fit	Brown (2006) Hu & Bentler (1999)
NFI	.94	NFI $\geq$ .95	NFI $\geq$ .90	Good fit	Tabachnick & Fidell (2007) Thompson (2008)
NNFI	.97	NNFI $\geq$ .95	NNFI $\geq$ .90	Perfect fit	Tabachnick & Fidell (2007) Thompson (2008)
CFI	.97	CFI $\geq$ .95	CFI $\geq$ .90	Perfect fit	Hu & Bentler (1999) Tabachnick & Fidell (2007)
GFI	.83	GFI $\geq$ .95	GFI $\geq$ .90	Poor fit	Hooper & colleagues (2008) Hu & Bentler (1999)

$\chi^2=1115.99$ ;  $df=551$

As can be seen in Table 2, it was found out that except for the GFI indices, all the other fit indices had perfect or good fit values. Thus, it could be stated that the model determined to have four factors was confirmed in Azerbaijani language. Table 3 demonstrates the correlations between each factor in the scale and the total-scale as well as those between the factors themselves.

Table 3. Pearson Correlation Coefficients between IAS and the Sub-Factors

Factor	Withdrawal	Controlling Difficulty	Disorder in Functionality	Social Isolation	Total-Scale
Withdrawal	1				
Controlling Difficulty	.631*	1			
Disorder in Functionality	.512*	.689*	1		
Social Isolation	.460*	.532*	.572*	1	
Total-Scale	.843*	.883*	.818*	.721*	1

\* $p < .001$

As can be seen in Table 3, significant correlations were found between each factor and the total-scale ( $p < .001$ ). In addition, significant correlations were also found between the sub-factors. The fact that there were correlations between the sub-factors proved that separate measurements could be conducted for each factor and that a single measurement could be done using the total scores regarding the total-scale.

### 3.1.3 Reliability Analysis Findings as a Result of CFA

As a result of CFA, the Cronbach's Alpha ( $\alpha$ ) internal consistency coefficient for the total-scale made up of four factors was calculated as  $\alpha = .926$ , while it was  $\alpha = .844$  for the factor of *Withdrawal*;  $\alpha = .821$  for *Controlling Difficulty*;  $\alpha = .829$  for *Disorder in Functionality*;  $\alpha = .792$  for *Social Isolation*.

## 3.4 Findings regarding the Prevalence of Internet Addiction in Azerbaijan

### 3.4.1 Examining the Assumptions

Regarding the data collected from 836 participants found in the sample. 4 students with outliers were excluded from the data set. As the values of both z-scores and Mahalanobis output for the remaining 832 students were in acceptable range, no other outliers were observed in the data set. The values of skewness (.391;  $\pm 1$ ) and kurtosis (.037;  $\pm 1$ ) were found to be in acceptable range. In addition, when the histogram, Kolmogorov-Smirnov Normality Test (.055;  $p > .05$ ), P-P and Q-Q graphics were examined, it was seen that the distributions were normal.

### 3.4.2 Reliability Analysis Findings as a Result of Prevalence

As a result of CFA, the Cronbach's Alpha ( $\alpha$ ) internal consistency coefficient for the total-scale made up of four factors was calculated as  $\alpha = .927$ , while it was  $\alpha = .836$  for the factor of *Withdrawal*;  $\alpha = .821$  for *Controlling Difficulty*;  $\alpha = .839$  for *Disorder in Functionality*;  $\alpha = .779$  for *Social Isolation*.

Table 4. Descriptive statistics regarding total-scale and sub-scale

Factor	N	Mean	Mean/Items	Sd.
Withdrawal	832	28,962	2,632	8,282
Controlling Difficulty	832	22,242	2,224	7,384
Disorder in Functionality	832	14,287	2,041	5,661
Social Isolation	832	12,208	1,744	4,753
Total-scale	832	77,698	2,220	21,656

Table 5. Grouping of the IAS Total Scores with Two-Step Cluster Analysis

IAS				
Group	N	%	Mean	Sd
1 (non-addicted group)	432	51.9	61,294	11,604
2 (addiction risk group)	338	40.6	90,249	8,880
3 (addicted group)	62	7.5	123,581	9,134
Total	832	100.0	77,698	21,656

When Table 5 is examined, it is seen that the number of Internet addicts was 62 (7.5%) and that the number of moderate Internet addicts (addiction risk group) was 338 (40.6%). The first group represented the non-addicted participants, and the second group represented the participants with the risk of addiction. Regarding this grouping, it could be stated that the first group did not have any symptoms of addiction at all; that the second group with the risk of addiction showed some of the symptoms of addiction, and that the third group, the addicted group, showed most of the symptoms of addiction.

In addition, the first group represented the non-addicted participants with Internet addictions, and the second group represented the participants with the risk of addiction. Regarding this grouping, it could be stated that the first group did not have any symptoms of addiction at all; that the second group with the risk of addiction showed some of the symptoms of addiction (some of items were 4 or 5 point), and that the third group, the addicted group, showed most (or all) of the symptoms of addiction.

#### 4. Conclusion and Discussion

In this study, "Internet Addiction Scale" developed by Gunuc & Kayri (2010), was adapted into the Azerbaijani language. In this sense, instead of developing a new measure, it was considered convenient based on cultural and language similarities to use a measure with validity and reliability proven in many studies. It attracts attention that no Internet addiction scale was developed and the prevalence of Internet addiction was not examined in Azerbaijan previously. As this study is the first in this specific area and would be a pioneer for future research, it is expected to significantly contribute, particularly, to the literature in Azerbaijan.

This research was conducted in two steps. In the first step, the validity and reliability studies for the scale were conducted. Upon having the factor structure of scale confirmed, the prevalence of Internet addiction was determined in a larger sample. In both steps, students were randomly selected from different majors, through stratified sampling. Data were obtained from 309 students in the sample of the first administration and 836 students in the sample of second administration.

The four-factor structure of the scale was tested through CFA. The Cronbach's Alpha ( $\alpha$ ) internal consistency coefficient of the scale was calculated as  $\alpha=.926$ . Reliability coefficient of 4 sub-factors, regarding the scale, was found as follows:  $\alpha=.844$  for *Withdrawal*,  $\alpha=.821$  for *Controlling Difficulty*,  $\alpha=.829$  for *Disorder in Functionality* and  $\alpha=.792$  for *Social Isolation*. Both item values and CFA indices were obtained on good levels. Both factor structure and item number of the original scale were preserved and confirmed. Based on these results, scale structure was shown to be robust enough to use in Azerbaijan.

Each item in the scale is positive and the total score obtained in the scale refers to the Internet addiction. In this sense, when the scale is used, comparison or relational analyses can be conducted between the total score obtained in the scale and the demographic variables. Besides, upon two-step cluster analysis over total scores, Internet addicts and non-addicts can be assigned into groups. Re-comparative analyses can be conducted in these newly drawn groups. In addition, Internet addicted persons can be scored through a quite simpler but scientifically weak method. In this sense, those over 105 (3\*35), as the total score obtained in the scale, can be considered Internet addicts. These can be categorized into three groups such as non-Internet addicts, moderate Internet addicts, and Internet addicts. Then, those between 105 (3\*35) and 140 (4\*35) may be considered moderate Internet addicts as those over 140 can be considered Internet addicts. In this context, moderate Internet addiction means that individuals have some symptoms of Internet

addiction and Internet addicts, on the other hand, have most or all symptoms of Internet addiction. It must be taken into consideration that all scoring methods, mentioned and not mentioned, can be used based on researchers' experiences, however, two-step cluster as a scientific method is recommended for stratifying individuals by the researchers in this study. In addition, comparing total scores to other variables and directly including them in relational analyses are also recommended as a scientific method.

In the second step, prevalence of Internet addiction among adolescents and youth in Azerbaijan was determined. It was found in the research that 51.9% were not Internet addicts, 40.6% were moderate Internet addicts, and 7.5% were Internet addicts. These findings show that the rate of Internet addicts, particularly with moderate Internet addicts and Internet addicts (48.1%), is not small enough to ignore. A comparison with other rates or different samples is not possible as there are no studies available in the national literature on the prevalence of Internet addiction in Azerbaijan. However, a review of findings in this context within international literature reveals similar rates in many countries. Yet, it is remarkable that moderate Internet addicts account for 40.6% in this research, compared to 26.4% in Turkey (Gunuc & Kayri, 2010) and 5.01% in Italy (Poli & Agrimi, 2012). Besides, it could be stated that these differences from one society to another stem from a number of factors such as culture, access to technology, age, parental education, level of income, different measurement tools and different measurement techniques, time of research conducted and selection of research samples (Durkee et al., 2012; Tsitsika et al., 2014).

Some precautions and treatments for addicts must be planned based on findings of future researches similar to this study in Azerbaijan. In order for moderate Internet addicts, in other words, users at risk, not to turn into addicts in time, families and educators must take on some responsibilities. However, prior to implementing any precautions or treatments, possible motives for Internet addiction must be defined particularly through case study method. Although international literature presents some findings on these factors, they may have different effect levels based on cultural differences. In this context, the Internet addiction scale adapted within this study is suggested for use as a diagnosis tool in clinical settings. In addition, investigating various demographic variables in different sample groups in Azerbaijan and evaluating the results following this process will present more robust data.

An important limitation within this study is that data were collected at one university. Even though, the number of samples is enough, a more comprehensive study to be conducted in Azerbaijan may more clearly reveal the prevalence of Internet addiction. The focus of this research has been about contributing a measurement tool needed in Azerbaijan in the national literature. Also, lack of prevalence of empirical studies in Azerbaijan prevented data collection from universities. With the introduction of measurement tool developed within this study, an increase in future empirical research is expected.

## References

- Brown, T. A. (2006). *Confirmatory factor analysis for applied research*. New York, NY: Guilford Publications, Inc.
- Byrne, B. (2010). *Structural equation modeling with AMOS: Basic concepts, applications and programming*. New York, NY: Taylor and Francis Group.
- Caplan, S. E. (2002). Problematic internet use and psychosocial well-being: development of a theory-based cognitive behavioral measurement instrument. *Computer Human Behavior, 18*, 553-575. [http://dx.doi.org/10.1016/S0747-5632\(02\)00004-3](http://dx.doi.org/10.1016/S0747-5632(02)00004-3)
- Caplan, S. E. (2010). Theory and measurement of generalized problematic internet use: a two-step approach. *Computer Human Behavior, 26*, 1089-1097. <http://dx.doi.org/10.1016/j.chb.2010.03.012>
- Chen, S., Weng, L., & Su, Y., et al. (2003). Development of a Chinese Internet addiction scale and its psychometric study. *Chinese Journal of Psychology, 45*, 279-294.
- Clark, D. J., & Frith, K. H. (2005). The development and initial testing of the Internet Consequences Scales (ICONS). *Computer Information Nurs, 23*, 285-291. <http://dx.doi.org/10.1097/00024665-200509000-00013>
- Demetrovics, Z., Szeredi, B., & Rózsa, S. (2008). The three-factor model of internet addiction: The development of the problematic internet use questionnaire. *Behavior Research Methods, 40*(2), 563-574. <http://dx.doi.org/10.3758/BRM.40.2.563>
- Durkee, T., Kaess, M., & Carli, V., et al. (2012). Prevalence of pathological internet use among adolescents in Europe: Demographic and social factors. *Addiction, 107*, 2210-2222. <http://dx.doi.org/10.1111/j.1360-0443.2012.03946.x>
- Everitt, B. S. (1980). *Cluster Analysis* (2<sup>nd</sup>ed.). London: *Heinemann Educational Books*.
- Gunuc, S. (2015). Relationships and Associations between Video Game and Internet Addictions: Is Tolerance a Symptom Seen in All Conditions? *Computer in Human Behavior, 49*, 517-525. <http://dx.doi.org/10.1016/j.chb.2015.03.063>



- Gunuc, S. (2013). Cart and Chaid analyses of some variables that predict internet addiction. *Turkish Journal of Psychology*, 28(71), 88-101.
- Gunuc, S., & Dogan, A. (2013). The Relationships between Turkish Adolescents' Internet Addiction, Their Perceived Social Support and Family Activities. *Computers in Human Behavior*, 29(6), 2197-2207. <http://dx.doi.org/10.1016/j.chb.2013.04.011>
- Gunuc, S. (2011). Examination of Internet Addiction among Computer Department Staff in Terms of Duration and Purpose of Its Use. *International Journal of Cyber Society and Education*, 4(1), 39-46.
- Gunuc, S., & Kayri, M. (2010). The profile of Internet Dependency in Turkey and Development of Internet Addiction Scale: Study of Validity & Reliability. *Hacettepe University Journal of Education*, 39, 220-232.
- Hambleton, R. K., & Patsula, L. (1999). Increasing the validity of adapted tests: myths to be avoided a guidelines for improving test adaptation practices. *Journal of Applied Testing Technology*, August 1999 Issue. <http://data.memberclicks.com/site/atpu/volume%201%20issue%20Increasing%20validity.pdf>
- Hooper, D., Coughlan, J., & Mullen, M. (2008). Structural equation modelling: guidelines for determining model fit. *Articles*, 2. <http://dx.doi.org/10.1007/978-3-319-19027-3>
- Hu, L. T., & Bentler, P. M. (1999). Cutoff Criteria for Fit Indexes in Covariance Structure Analysis: Conventional Criteria Versus New Alternatives. *Structural Equation Modeling*, 6(1), 1-55. <http://dx.doi.org/10.1080/10705519909540118>
- Huck, S. (2012). *Reading statistics and research* (6<sup>th</sup> ed.). Boston: Pearson.
- Hutcheson, G. D., & Sofroniou, N. (1999). *The multivariate social scientist: An introduction to generalized linear models*. Sage Publications.
- Jöreskog, K., & Sörbom, D. (2001). *Lisrel 8: User's reference guide*. Chicago, USA: Scientific Software International Inc.
- Kaufman, L., & Rousseeuw, P. J. (1990). Finding groups in data: An introduction to cluster analysis. New York: John Wiley and Sons. <http://dx.doi.org/10.1002/9780470316801>
- Kim, K., Ryu, E., & Chon, M. Y., et al. (2006). Internet addiction in Korean adolescents and its relation to depression and suicidal ideation: a questionnaire survey. *International Nurs Stud*, 43, 185-192. <http://dx.doi.org/10.1016/j.ijnurstu.2005.02.005>
- Kline, R. B. (2011). *Principles and practice of structural equation modeling* (3th ed.) New York: The Guilford Press.
- Morahan-Martin, J., & Schumacher, P. (2000). Incidence and correlates of pathological internet use among college students. *Computers in Human Behavior*, 16, 13-29. [http://dx.doi.org/10.1016/S0747-5632\(99\)00049-7](http://dx.doi.org/10.1016/S0747-5632(99)00049-7)
- Nichols, L. A., & Nicki, R. (2004). Development of a Psychometrically Sound Internet Addiction Scale: A Preliminary Step. *Psychology of Addictive Behaviors*, 18(4), 381-384. <http://dx.doi.org/10.1037/0893-164X.18.4.381>
- Pallant, J. (2007). *SPSS survival manual: A step by step guide to data analysis using SPSS for Windows*. Maidenhead: Open University Press.
- Poli, R., & Agrimi, E. (2012). Internet addiction disorder: Prevalence in an Italian student population. *Nordic Journal of Psychiatry*, 66(1), 55-59. <http://dx.doi.org/10.3109/08039488.2011.605169>
- Raykov, T., & Marcoulides, G. A. (2006). *A first course in structural equation modeling* (2nd ed. Lawrence Erlbaum Associates, Inc. Publishers.
- Tabachnick, G. G., & Fidell, L. S. (2007). *Experimental designs using ANOVA*. Belmont, CA: Duxbury, 35, 793-809.
- Thatcher, A., & Goolam, S. (2005). Development and psychometric properties of the Problematic Internet Use Questionnaire. *South African Journal Psychology*, 35, 793-809. <http://dx.doi.org/10.1177/008124630503500410>
- Tsitsika, A., Janikian, M., & Schoenmakers, T. M., et al. (2014). Internet Addictive Behavior in Adolescence: A Cross-Sectional Study in Seven European Countries. *Cyberpsychology, Behavior, and Social Networking*, 17(8), 528-535. <http://dx.doi.org/10.1089/cyber.2013.0382>
- Wölfling, K., & Müller, K. W. (2010). Pathological gambling and computer game-addiction. Current state of research regarding two subtypes of behavioral addiction. *Bundesgesundheitsblatt, Gesundheitsforschung, Gesundheitsschutz*, 53(4), 306-312. <http://dx.doi.org/10.1007/s00103-010-1038-7>
- Wu, X., Chen, X., Han, J., Meng, H., & Luo, J., et al. (2013). Prevalence and Factors of Addictive Internet Use among Adolescents in Wuhan, China: Interactions of Parental Relationship with Age and Hyperactivity-Impulsivity. *PLoS*

ONE 8(4): e61782. <http://dx.doi.org/10.1371/journal.pone.0061782>

Young, K. S. (1998). Internet Addiction: The Emergence of a New Clinical Disorder. *CyberPsychology and Behavior*, 1(3), 237-244. <http://dx.doi.org/10.1089/cpb.1998.1.237>

Zboralski, K., Orzechowska, A., & Talarowska, M., et al. (2009). The prevalence of computer and Internet addiction among pupils. *Postepy Higieny I Medycyny Do wiazczalnej* (Online), 63, 8–12.

Statistics Azerbaijan. (2010). Retrieved from [http://www.stat.gov.az/menu/6/statistical\\_yearbooks/](http://www.stat.gov.az/menu/6/statistical_yearbooks/)

Internet World Stats. (2015). Retrieved from <http://www.internetworldstats.com/stats.htm>

#### Appendix. Internet Addiction Scale for Azerbaijani Form

<b>İNTERNETDƏN ASILILIQ TESTİ</b> Aşağıdakı hər bir maddəni oxuyun, bu maddə sizing üçün hər zaman doğru isə “tamamilə razıyam”, ümumilikdə doğru isə “razıyam”, əmin deyilsinizsə, “qərarızsam”, ümumilikdə razı deyilsinizsə “razı deyiləm”, heç vaxt doğru deyilsə, “qəti razı deyiləm” cavablarını işarələməyiniz xahiş olunur.	Qəti razı deyiləm.	Razı deyiləm	Qərarızsam	Razıyam	Tamamilə razıyam
<b>Əksiklik</b>					
1. İnternetdən istifadə etmədiyim zaman özümü gərgin, narahat hiss edirəm.					
2. İnternetdən istifadə etmək istəyib də edə bilmədikdə çox hirsli və qəzəbli oluram.					
3. İnternet əlaqəsi kəsildə ya da yavaşlayanda hirsələnirəm / qəzəblənirəm.					
4. İnternetdən istifadə etdiyim zaman heç vaxt olmadığım qədər özümü xoşbəxt hiss edirəm.					
5. Özümü narahat və sıxıntılı hissetdiyim zamanlarda internetdən istifadə etmək məni sakitləşdirir.					
6. Kimsə məni internetdən ayırırsa hirsələnirəm.					
7. Problemlərimdən qaçmaq üçün internetdən istifadə edirəm.					
8. İnternetə planlaşdırdığım vaxt girmədiyimdə hirsələnirəm.					
9. Ətrafımda insanlar olduğu vaxt tək qalıb internetə girmək istəyirəm.					
10. İnternetdən istifadə etmədiyim zaman internetə girmək üçün səbirsizlənir və tələsirəm.					
11. Getdiyim məkanlarda internet əlaqəsi axtarıram.					
<b>Nəzarət etmədə çətinlik</b>					
12. İnternet istifadəsinə məhdudiyət qoymaqda və nəzarət etməkdə çətinlik çəkirəm.					
13. Səhər oyananda ilk ağılıma gələn fikir internetə girməkdir.					
14. İnternetdə hər səfərimdə əvvəlkindən daha uzun müddət qalmaq istəyirəm.					
15. İnternetdə planlaşdırdığımdan daha uzun müddət qalırım.					
16. İnternetdən istifadə etmədiyim zamanlarda belə internet fikirləşirəm.					
17. İnternetdə ikinə aqlığı, susuzluğumu hiss etmirəm ya da fərqinə varmıram.					
18. İnternetdə daha uzun vaxt keçirmək üçün başqa planlarımı ləğvedirəm.					
19. İstədiyim zaman internetdən ayrılı bilmirəm.					
20. Ailəm məni çağırırsa da internetdən ayrılı bilmirəm.					
21. İnternetdən istifadə etmək üçün yuxumu ertələyirəm.					
<b>Funksional pozulma</b>					
22. İnternetdən istifadə etdiyim üçün ailəmlə problemlər yaşayıram.					
23. Dostlar məni çağırırsa da internetdən ayrılı bilmirəm.					
24. İnternet istifadə etdiyim üçün başqa fəaliyyətlərə marağım azalır.					
25. İnternet istifadə etdiyimə görə ev/iş/məktəb məsuliyyətlərimi yerinə yetirə bilmirəm ya da etmək istəmirəm.					
26. Ətrafımdakilər internetdə sərfələdiyim zamana görə şikayət edirlər.					
27. İnternetdən istifadə etdiyim üçün ailəmlə daha az zaman keçirirəm.					
28. İnternetdən istifadə etdiyim üçün dostlarım ilə daha az zaman keçirirəm.					
<b>Sosial təcridolma</b>					
29. İnternetdən istifadə etdiyim üçün yoldaşlarım ilə problemlər yaşayıram.					
30. Real həyatdakı dostluqlarımdansa İnternet mühitində olan dostluqlarımı seçərdim.					
31. Real həyatdakı dostlarımla çöldə görüşmək yerinə internetdə görüşməyə üstünlük verərəm.					
32. Dostlarımı internet üzərindən tapıram.					
33. İnternet mənim ən yaxşı dostumdur.					
34. İnternetsiz həyat mənə mənasız və boş gəlir.					
35. İnternet istifadə etdiyim üçün üz üzə ünsiyyət qurmaqda çətinlik çəkirəm.					



This work is licensed under a [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/).